

Revised in August 2010

Please read this insert before taking a drug.
Further, keep this insert so that you can read
it whenever necessary.

Living bacteria preparation for intestinal regulation

Designated Quasi-Drug

Bio-Three[®] Hi

Features:

Bio-Three Hi is a living bacteria preparation containing 3 different kinds useful bacteria such as Lactic acid bacteria, Butyric acid bacteria and Amyolytic bacteria. By active proliferation of these 3 kinds useful bacteria and the incidental increase of bifidobacteria, intestinal microflora becomes well balanced, consequently fecal disorder improves including constipation and loose stool.

Caution in use:

1. Following persons should consult with physician or pharmacist before taking.
Those persons who are currently receiving medical treatment by physician.
2. In the following case, discontinue taking and consult with physician or pharmacist by submitting this insert.

When symptoms are not improved after taking for about 1 month.

Ingredients & Content (per 6 tablets):

| Ingredient | Content | Action |
|------------------------------------|---------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Lactomin (Lactic acid bacteria) | 30mg | It produces lactic acid which regulates intestinal environments, and improves stool disorders by symbiosis with butyric acid bacteria. |
| Butyric acid bacteria | 150mg | It proliferates in the intestine with lactic acid bacteria and produces butyric acid which is useful for intestinal health, and activates intestinal movements. |
| Amyolytic acid bacteria | 150mg | It proliferates bifidobacteria and lactic acid bacteria which are useful intestinal bacteria. |

As additives, Polyvinyl-alcohol (Fully hydrolized), Povidone, Potato-starch and Lactose hydrate are contained.

Indications:

Intestinal regulation (fecal regulation), Constipation, Soft stool, Abdominal fullness



Those with constipation



Those with soft stool



Those with abdominal fullness

Dosage and administration:

Take the followings after each meal.

| Age | Each dosage | Daily frequency |
|-----------------------|-------------------|-----------------|
| | tablet | |
| Adult (more than 15y) | 2 tablets | 3 times |
| Infant (3m – 15y) | 1 tablet | |
| Baby (less than 3m) | No administration | |

Caution for dosage & administration:

For infant's administration, parents should take care of them.

Caution for storage and handling:

- (1) Keep with sealing in a dry place avoiding direct sunlight.
- (2) Keep away from children.
- (3) Do not change to other container. (To prevent misuse and deterioration)
- (4) Bottle package: Tighten the cap firmly every time after taking the tablet because it is easily affected by moisture.
- (5) Bottle package: Remove stuffing after opening the cap. (Packing material in the upper part of bottle is provided to prevent the breaking of tablets.)
- (6) Do not use after expiration.

Packing: 270 tablets/bottle

For inquiry: For any inquiry about this product, please contact the below:

Domestic distributor:

TOA Shinyaku Co. Ltd.

3-2-11 Nishi-Shinjuku, Shinjuku-Ku, Tokyo

Tel: 03-3347-0770, Fax: 03-3347-0780

9:00am – 17:00pm (Week days only)

<http://www.toashinyaku.co.jp/>

Manufacturer: TOA Pharmaceutical Co., Ltd. 2-1-11 Sasazuka, Shibuya-Ku, Tokyo

Domestic distributor: TOA Shinyaku Co. Ltd. 3-2-11 Nishi-Shinjuku, Shinjuku-Ku, Tokyo